## Nontechnical Abstract

The purpose of this clinical investigation is to determine if the drug pVGI.1(VEGF2) is safe and effective in promoting the development of new blood vessels in the heart. This may increase blood flow to the heart and consequently reduce chest pain and/or increase exercise tolerance in patients with myocardial ischemia.

Myocardial ischemia is a condition in which there is inadequate blood flow to the heart resulting in chest pain. Often myocardial ischemia is caused by coronary artery disease (CAD). Cardiovascular disease is the number one cause of mortality in the United States, and most cardiovascular deaths are due to CAD. While various medications, angioplasty and/or coronary bypass surgery may be sufficient to relieve chest pain in some patients, these methods are not suitable options for others. Therefore, Vascular Genetics Inc. is investigating the use of gene therapy as a treatment for increasing blood flow to the heart in patients with no other therapeutic options.

The treatment will involve using a catheter to deliver deoxyribonucleic acid (DNA), or genetic material, into the heart. Once inside the heart muscle, this DNA will direct the muscle cells to make a specific protein called vascular endothelial growth factor 2 (VEGF-2). VEGF-2 is a protein that has been shown to cause the growth of new blood vessels under a variety of conditions. Experiments in the laboratory have suggested that VEGF-2 gene therapy can be used to grow new blood vessels in animals whose arteries have been surgically blocked. Additionally, VEGF-2 gene therapy is being used to increase blood flow and reduce resting leg pain (i.e., ischemic rest pain) in the feet or legs of patients with critical limb ischemia, or leg pain due to decreased blood flow in legs due to blocked arteries.

This study will last up to 13 months (which includes a Pretreatment Phase, a Treatment Phase, a Post-treatment phase, and a Follow-up Phase). Eighteen to 27 patients will be enrolled and divided into three groups; each group will receive one of three doses of pVGI.1(VEGF2). Two patients will receive pVGI.1(VEGF2) to every one patient who receives placebo. Patients who have not improved will have the opportunity to enroll in another study to receive pVGI.1(VEGF2) treatment after completing the initial Post-treatment Phase of this study (3 months).

In summary, Vascular Genetics Inc. is investigating the possibility that the delivery of VEGF-2 DNA into the oxygendeprived heart will result in the development of new blood vessels and thereby increase the blood supply to the heart and reduce chest pain.